

09/674,819

-15-

PD-5774-01-DRK

**REMARKS**

Claims 1-17 are pending in the application. Only Claims 1 and 12 are in independent form. New Claims 18 and 19 have been added by way of the present amendment.

The application has been held to contain claims directed to more than one species of generic invention thereby being deemed to lack unity of invention as to not form a single general inventive concept under PCT Rule 13.1. The claims were held to contain a number of species for the active compound, the humectants, and optional auxiliary agents, to which the claims shall be restricted if no claim is finally held to be allowable. Applicant was required, in reply to the Office Action, to elect a single species from each of the active compounds, the humectants, and the optional auxiliary agents, to which the claims shall be restricted if no generic claim is finally held to be allowable. Applicant hereby provisionally elects gabapentin as the single species for the active compounds and propylene glycol as the single species for the humectants. Applicant has amended the claims to remove the auxiliary agent from the independent claims thereby eliminating the need to make a species election therefor. The claims readable on the elected species include Claims 1, 2, 4 and 8-19.

Applicant respectfully traverses the lack of unity of invention under Rule 13.1 as the active compounds all share the general formula as set forth in Claim 1 in common. Accordingly, the Applicant respectfully submits that the lack of unity rejection under PCT Rule 13.1 is not applicable to the present claims and Applicant respectfully submits that all species set forth in the application should be examined as a single general inventive concept under PCT Rule 13.1.

However, as set forth in the Office Action, Applicant has made the required species elections.

In view of the foregoing amendments and remarks, Applicants respectfully submits that the application is in condition for allowance, which allowance is respectfully requested.

09/674,819

-16-

PD-5774-01-DRK

The Commissioner is authorized to charge any fee or credit any overpayment in connection with this communication to our Deposit Account 23-0455.

Respectfully submitted,




David R. Kurlandsky  
Reg. No. 41,505  
Warner-Lambert Company  
2800 Plymouth Road  
Ann Arbor, MI 48105  
Tel. (734) 622-5215  
Fax (734) 622-1553

Attachment - Amended claims, Version with Markings to Show Changes Made

DK1P4444.doc

**"VERSION WITH MARKINGS TO SHOW CHANGES MADE"****IN THE CLAIMS:**

Claim 1 (amended). A stabilized solid composition containing a 4-amino-3-substituted-butanoic acid derivative which comprises a 4-amino-3-substituted-butanoic acid derivative having the general formula



wherein,

R<sub>1</sub> is a hydrogen atom, a hydroxyl group, a methyl group or an ethyl group;

R<sub>2</sub> is a monovalent group selected from:

a straight or branched alkyl group of 3 - 8 carbon atoms;

a straight or branched alkylene group of 3 - 8 carbon atoms;

a straight or branched alkyl group of 3 - 8 carbon atoms which is mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 3 - 8 carbon atoms;

a cycloalkyl group of 3 - 8 carbon atoms which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms;

09/674,819

-18-

PD-5774-01-DRK

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms, is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-, and one or two of the unsubstituted methylene groups (-CH<sub>2</sub>-) are mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-;

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-, and one or two of the unsubstituted methylene groups (-CH<sub>2</sub>-) being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-, said phenyl group being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, and one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>- and one or two of the unsubstituted methylene groups (-CH<sub>2</sub>-) being mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl

group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a phenyl or naphthyl group;

a phenyl group substituted with a methylenedioxy group;

a phenyl or naphthyl group which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group, a phenoxy group, a phenylmethoxy group, a phenylmethoxy group wherein said phenyl ring is mono-substituted with a halogen atom, trifluoromethyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring, a cycloalkanedienylmethoxy group having 5 - 8 carbon atoms in the cycloalkanedienyl ring, a cycloalkylmethoxy group wherein one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-, a cycloalkenylmethoxy group wherein one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-, a cycloalkanedienyl-methoxy group wherein one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkanedienyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>- group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring wherein said cycloalkyl ring is mono-substituted with a halogen atom, trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring wherein said cycloalkenyl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-, or a cycloalkanedienylmethoxy group having 5 - 8 carbon atoms in the cycloalkanedienyl ring wherein said cycloalkanedienyl ring is mono-substituted with a halogen atom, a

09/674,819

-21-

PD-5774-01-DRK

trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkanediényl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-;

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS-;

an -O-, -S- or -SS-phenyl group;

a diphenylamino group;

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, a alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an -O-, -S- or -SS-phenyl group wherein said phenyl group is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

or

R<sub>1</sub> and R<sub>2</sub>, together with the carbon atom to which they are attached, may form a divalent group selected from:

a cycloalkylidene group of 5 - 8 carbon atoms;

a cycloalkylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group or a carboxyl group;

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-;

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>- group and one or more of the unsubstituted methylene groups (-CH<sub>2</sub>-) in said cycloalkyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>- group and one or more of the unsubstituted methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring or cycloalkanedienyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms, said phenyl ring being mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl



09/674,819

-23-

PD-5774-01-DRK

group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;


a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group; and

a humectant[; and, if necessary, an auxiliary agent] for manufacturing a pharmaceutical preparation.

Claim 9 (amended). The stabilized solid composition containing a 4-amino-3-substituted-butanoic acid derivative as claimed in claim [1] 18 wherein a total amount of said humectant is 0.01 - 25% by weight relative to a total amount of the 4-amino-3-substituted-butanoic acid derivative and an auxiliary agent for manufacturing a pharmaceutical preparation.

Claim 12 (amended). A process for the preparation of a solid composition containing a 4-amino-3-substituted-butanoic acid derivative having the general formula



wherein,

R<sub>1</sub> is a hydrogen atom, a hydroxyl group, a methyl group or an ethyl group;

R<sub>2</sub> is a monovalent group selected from:

a straight or branched alkyl group of 3 - 8 carbon atoms;

a straight or branched alkylene group of 3-8 carbon atoms;

a straight or branched alkyl group of 3 - 8 carbon atoms which is mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkoxy

group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 3 - 8 carbon atoms;

a cycloalkyl group of 3 - 8 carbon atoms which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedieryl group of 5 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedieryl group of 5 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms, is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-;

09/674,819

-25-

PD-5774-01-DRK

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-, and one or two of the unsubstituted methylene groups (-CH<sub>2</sub>-) are mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedieryl group of 5 - 8 carbon atoms, one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring or cycloalkanedieryl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-;

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedieryl group of 5 - 8 carbon atoms, one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring or cycloalkanedieryl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-, and one or two of the unsubstituted methylene groups (-CH<sub>2</sub>-) being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-, said phenyl group being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedieryl group of 5 - 8 carbon atoms, one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring or cycloalkanedieryl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedieryl group of 5 - 8 carbon atoms,

one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, and one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>- and one or two of the unsubstituted methylene groups (-CH<sub>2</sub>-) being mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a phenyl or naphthyl group;

a phenyl group substituted with a methylenedioxy group;

a phenyl or naphthyl group which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group, a phenoxy group, a phenylmethoxy group, a phenylmethoxy group wherein said phenyl ring is mono-substituted with a halogen atom, trifluoromethyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring, a cycloalkanedienylmethoxy group having 5 - 8 carbon atoms in the cycloalkanedienyl ring, a cycloalkylmethoxy group wherein one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-, a cycloalkenylmethoxy group wherein one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or

09/674,819

-27-

PD-5774-01-DRK

-S(O)<sub>2</sub>-, a cycloalkanedieryl-methoxy group wherein one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkanedieryl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>- group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring wherein said cycloalkyl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring wherein said cycloalkenyl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-, or a cycloalkanedieryl-methoxy group having 5 - 8 carbon atoms in the cycloalkanedieryl ring wherein said cycloalkanedieryl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkanedieryl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-;

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS-;

an -O-, -S- or -SS-phenyl group;

a diphenylamino group;

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, a alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS- and mono-, di- or tri-substituted

09/674,819

-28-

PD-5774-01-DRK

with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an -O-, -S- or -SS-phenyl group wherein said phenyl group is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

or

R<sub>1</sub> and R<sub>2</sub>, together with the carbon atom to which they are attached, may form a divalent group selected from:

a cycloalkylidene group of 5 - 8 carbon atoms;

a cycloalkylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group or a carboxyl group;

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>-;

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)<sub>2</sub>- group and one or more of the unsubstituted methylene groups (-CH<sub>2</sub>-) in said cycloalkyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedierylidene group of 5 - 8 carbon atoms;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedierylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedierylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) in said

09/674,819

-29-

PD-5774-01-DRK

cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>-;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)<sub>2</sub>- group and one or more of the unsubstituted methylene groups (-CH<sub>2</sub>-) in said cycloalkenyl ring or cycloalkanedienyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms, said phenyl ring being mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group, which comprises combining the 4-amino-3-substituted-butanoic acid derivative with a humectant [and, if necessary, an auxiliary agent] for manufacturing a pharmaceutical preparation.

Add the following new Claims 18 and 19.

Claim 18 (new). The stabilized solid composition containing a 4-amino-3-substituted butanoic acid derivatives as claimed in Claim 1 is further combined with an auxiliary agent.

09/674,819

-30-

PD-5774-01-DRK

Claim 19 (new). The process as claimed in Claim 12 wherein the solid composition further includes an auxiliary agent.